

The effective ω -meson mass with a vector field self-interaction

Katsuaki Sakamoto, Hiroaki Kouno and Akira Hasegawa
Department of Physics, Saga University, Saga 840, Japan

and

Masahiro Nakano
University of Occupational and Environmental Health, Kitakyushu 807, Japan

Abstract

We calculate the effective mass m_ω^* of the ω -meson in nuclear medium by the σ - ω model with a vector field self-interaction (VFSI). We found that the direct and the indirect effects of the VFSI almost cancel each other and the result is changed only slightly at the normal density. The VFSI makes m_ω^* smaller at lower density and makes m_ω^* larger at higher density. However, both effects are not large.

The effective meson mass in nuclear medium is one of the most interesting issues in intermediate energy physics. Much attention has been paid to predicting the meson masses in nuclear medium from various approaches such as quantum hadrodynamics (QHD) [1]-[7], the QCD sum rules [8, 9] and the quark meson coupling model [10]. It is widely accepted that the effective mass of the vector meson (ω and ρ) decreases as density increases. However, quantitatively, the result is considerably variable in each model.

In QHD, there are some versions of the Lagrangian to reproduce the properties of the nuclear matter and finite nuclei. One interesting version of them is the σ - ω model with the vector field self-interaction (VFSI). [11] From phenomenological analyses, the VFSI is needed for a better fit of equation of state at high density. (See, e.g., reference [12].)

As usual, this VFSI is regarded as an effective interaction caused by the quantum corrections. This means that the effective potential of the VFSI is added to the effective potential of the total system. On the other hand, the effective meson mass can be defined as a second derivative of the effective potential with respect to the meson-field expectation value. Therefore, the VFSI just shifts the effective ω -meson mass. Naively the VFSI is expected to make the effective ω -meson mass larger, since the potential of the VFSI has a plus sign. Is it true or not? In this brief report,

we study the effective ω -meson mass in nuclear medium by the σ - ω model with the VFSI to answer this question.

We start from the Lagrangian of the σ - ω model with the meson self-interaction:

$$\begin{aligned} L &= \bar{\psi}(i\gamma_\mu\partial^\mu - M + g_\sigma\phi - g_\omega\gamma_\mu V^\mu)\psi + \frac{1}{2}\partial_\mu\phi\partial^\mu\phi - \frac{1}{2}m_\sigma^2\phi^2 - \frac{1}{4}F_{\mu\nu}F^{\mu\nu} + U_\omega + \delta L, \\ U_\omega &= \frac{1}{2}m_\omega^2 V_\mu V^\mu \left(1 + \frac{g_\omega^2}{2}Y^2 V_\mu V^\mu\right), \end{aligned} \quad (1)$$

where ψ , ϕ , V_μ , M , m_σ , m_ω , g_σ and g_ω are the nucleon field, the σ -meson field, the ω -meson field, the nucleon mass, the σ -meson mass, the ω -meson mass, the σ -nucleon coupling and the ω -nucleon coupling, respectively. The vector field strength is given by $F_{\mu\nu} = \partial_\mu V_\nu - \partial_\nu V_\mu$, and the constant parameter Y represents the strength of the ω -meson self-interaction. The δL is a counter term adjusted to satisfy renormalization conditions as in the ordinary relativistic Hartree approximation (RHA) [13]. We regard the VFSI as an effective interaction which is originated in the higher-order quantum corrections beyond the 1-loop approximation, namely RHA.

In the RHA, the meson fields are replaced by their ground-state expectation values, $\phi \rightarrow \phi_0$, $V_\mu \rightarrow \delta_{\mu 0}V_0$. The expectation value of the ω field is determined by the baryon density ρ , [11]

$$W = \frac{g_\omega^2}{m_\omega^2} \frac{\rho}{1 + Y^2 W^2} = \frac{g_\omega^2}{m_\omega^2} \frac{\rho}{1 + \frac{y^2 W^2 m_\omega^4}{g_\omega^4 \rho_0^2}}, \quad W = g_\omega V_0, \quad (2)$$

where $y = g_\omega^2 \rho_0 / m_\omega^2$ is a dimensionless parameter and ρ_0 is a saturation baryon density. The expectation value ϕ_0 is determined by equation of motion of the σ -meson. Namely,

$$g_\sigma \rho_s = m_\sigma \phi_0, \quad (3)$$

where ρ_s is the scalar density of nucleons. We remark that the vacuum fluctuation effects is already included in ρ_s in Eq. (3). The ϕ_0 shifts the nucleon mass from M to the effective mass M^* in nuclear medium as

$$M^* = M - g_\sigma \phi_0. \quad (4)$$

After y is given, the parameters $C_\sigma = g_\sigma M / m_\sigma$, $C_\omega = g_\omega M / m_\omega$ are chosen to reproduce the saturation conditions at the baryon density 0.15 fm^{-3} and at the binding energy -15.75 MeV . These two parameters, the effective nucleon mass M_0^* at saturation density and the incompressibility K are listed in Table 1.

The second order variation of the Lagrangian (1) around the classical value V_0 gives the inverse of the ω -meson propagator,

$$\left(\partial^\mu \partial^\rho - g^{\mu\rho} \partial_\sigma \partial^\sigma - \frac{\partial^2 U_\omega}{\partial V_\mu \partial V_\rho} \Big|_{\delta_{\mu 0} V_0} \right) D(x-y)_{\rho\nu} = -\delta_\nu^\mu \delta^{(4)}(x-y). \quad (5)$$

By taking Fourier transform, we obtain the inverse of the meson propagator in momentum space,

$$[D^{-1}(k)]_{\mu\nu} = [k^2 - m_\omega^2(1 + Y^2 W^2) + i\epsilon]g_{\mu\nu} - k_\mu k_\nu - 2m_\omega^2 Y^2 W^2 \delta_{\mu 0} \delta_{\nu 0}. \quad (6)$$

Model	y	M_0^*/M	K [MeV]	C_ω^2	C_σ^2
RHA	0	0.730	454.1	146.86	226.94
VS1	0.2	0.721	422.5	159.13	239.84
VS2	0.5	0.692	363.4	203.59	284.73

Table 1: The parameters in σ - ω model.

In order to compute the ω -meson full propagator in the nuclear medium we solve Dyson's equation in the random phase approximation (RPA). Dyson's equation becomes a matrix equation,

$$D'_{\mu\nu} = D_{\mu\nu} + D_{\mu\rho}\Pi^{\rho\sigma}D'_{\sigma\nu}. \quad (7)$$

We have omitted the σ - ω mixing components in Eq. (7), since we are interested only in the meson self-energy with vanishing external spatial momentum as is seen below. In the one-loop level, the RPA self-energy of the ω -meson is given by

$$\Pi_{\mu\nu}(q) = -ig_\omega^2 \int \frac{d^4k}{(2\pi)^4} \text{Tr}[\gamma_\mu G(k) \gamma_\nu G(k+q)], \quad (8)$$

where the nucleon propagator $G(k)$ in nuclear medium is given by

$$\begin{aligned} G(k) &= (\gamma^\mu k_\mu^* + M^*) \left[\frac{1}{k^{*2} - M^{*2} + i\epsilon} + \frac{i\pi}{E^*(k)} \delta(k_0^* - E^*(k)) \theta(k_F - |\mathbf{k}|) \right] \\ &\equiv G_F(k) + G_D(k), \end{aligned} \quad (9)$$

where k_F is the Fermi momentum, $k^{*\mu} = (k^0 - g_\omega V^0, \mathbf{k})$ and $E^*(k) = \sqrt{\mathbf{k}^2 + M^{*2}}$. We remark that the VFSI does not appear in RPA, since we regard it as an effective interaction induced by the higher-order quantum correction beyond the 1-loop approximation.

The equation (8) is divided into two parts

$$\Pi_{\mu\nu} = \Pi_{\mu\nu}^{\text{Den}} + \Pi_{\mu\nu}^{\text{Vac}} = \Pi_{\mu\nu}^{\text{Den}} + \left(g^{\mu\nu} - \frac{q^\mu q^\nu}{q^2} \right) \Pi^{\text{Vac}}. \quad (10)$$

The first term describes particle-hole excitation and the Pauli blocking for N and $\bar{N}$ excitation below the Fermi surface. The second term describes N and $\bar{N}$ excitation, this part generally has divergence and we remove the divergence by the renormalization procedure as in [1]. Namely,

$$\begin{aligned} \Pi^{\text{Vac}} \Big|_{q^2=m_\omega^2, \rho=0} &= 0, \\ \frac{\partial}{\partial q^2} \Pi^{\text{Vac}} \Big|_{q^2=m_\omega^2, \rho=0} &= 0. \end{aligned} \quad (11)$$

The effective meson mass in the nuclear medium is defined as the pole of the meson propagator Eq. (7), namely, given by the following equation,

$$m_\omega^{*2} = m_\omega^2 (1 + Y^2 W^2) + \Pi_T(q_0 = m_\omega^*, \mathbf{q} = 0), \quad (12)$$

where Π_T is the transverse component of ω -meson self-energy. Of course, the term including the parameter Y is occurred from the existence of the VFSI.

First, we examine only the direct effect on the effective ω meson by adding the VFSI to the effective potential. We use the effective ω meson mass which is defined by

$$m_\omega^{*2} = m_\omega^2(1 + Y^2 W^2). \quad (13)$$

In Fig. 1, we show the density dependence of m_ω^* with Eq. (13). The bold solid and bold dashed lines correspond to the results with the parameter sets in $y = 0.2$ and 0.5 , respectively. The m_ω^* increases as ρ increases. The effective potential for the VFSI increases m_ω^* largely as is expected.

Next, we study the VFSI effect on the effective ω -meson mass via the change of C_ω and M^* in Π_T . In this case, we do not contain the direct effect of the VFSI and the effective ω meson mass is defined by

$$m_\omega^{*2} = m_\omega^2 + \Pi_T(q_0 = m_\omega^*, \mathbf{q} = 0; C_\omega, M^*). \quad (14)$$

The numerical results of m_ω^* are also shown in Fig. 1. The solid, dashed and dotted lines correspond to the results with the parameter sets for RHA, VS1 and VS2 in table 1, respectively. This "indirect VFSI effect" via the change of C_ω and M^* in Π_T makes m_ω^* smaller, since the VFSI makes C_ω larger and makes M_0^* smaller.

Fig. 1

For a complete calculation in RPA with the VFSI we must use the Eq. (12). In Fig. 2, we show the density dependence of m_ω^* with the Eq. (12). We see that the direct and the indirect effects of the VFSI almost cancel each other at the normal density. The VFSI makes m_ω^* smaller at lower density via the large C_ω . The VFSI makes m_ω^* larger at higher density due to the direct effect of the VFSI. However, both effects are not large.

Fig. 2

In summary, we have studied the effect of the VFSI for the effective ω -meson mass in nuclear medium by using the RHA and RPA. In the calculation of the RHA, the VFSI increases C_ω to satisfy the saturation conditions. It also makes the values of M_0^* and the incompressibility K decrease. For the effective ω -meson mass in RPA, there is a large direct effect by adding the VFSI to the effective potential. However, the direct effect is almost canceled by the indirect effect via the change of M_0^* and C_ω in RPA calculations. As a result, the value of the effective ω -meson mass is changed only slightly at the normal density. The VFSI makes m_ω^* smaller at lower density and makes m_ω^* larger at higher density. However, both effects are not large.

Acknowledgements

The authors thank T. Kohmura and T. Maruyama for useful discussions.

References

- [1] H.-C. Jean, J. Piekarewicz, and A. G. Williams, Phys. Rev. **C49** (1994), 1981.
- [2] D. K. Griegel and T. D. Cohen, Phys. Rev. **C39** (1989), 1032.
- [3] H. Shiomi and T. Hatsuda, Phys. Lett. **B334** (1994), 281.
- [4] H. Kurasawa and T. Suzuki, Prog. Theor. Phys. **84**, (1990), 1030.
- [5] K. Tanaka, W. Bentz, A. Arima, and F. Beck, Nucl. Phys. **A528** (1991), 676.
- [6] M. Nakano, N. Noda, T. Mitsumori, K. Koide, H. Kouno and A. Hasegawa, Phys. Rev. **C55** (1997), 890.
- [7] K. Sakamoto, M. Nakai, H. Kouno, A. Hasegawa and M. Nakano, Int. J. Mod. Phys. **E9** (2000), 169.
- [8] T. Hatsuda and S. H. Lee, Phys. Rev. **C46** (1992), R34.
- [9] M. Asakawa and C. M. Ko, Phys. Rev. **C48** (1993), R526.
- [10] K. Saito and A. W. Thomas, Phys. Rev. **C51** (1995), 2757. ; K. Saito K. Tsushima and A. W. Thomas, Phys. Rev. **C55** (1997), 2637.
- [11] A. R. Bodmer, Nucl. Phys. **A526** (1991), 703.
- [12] Y. Sugahara and H. Toki, Nucl. Phys. **A579** (1994), 557.
- [13] S. A. Chin, Phys. Lett. **B62** (1976), 263; S. A. Chin, Ann. of Phys. **108** (1977), 301.




Figure 1: The effective ω -meson mass m_ω^* is shown as a function of the baryon density ρ . The bold solid and bold dashed lines correspond to the results for the direct effect with the parameter sets in $y = 0.2$ and 0.5 , respectively. The solid, dashed and dotted lines correspond to the results for the indirect effect with the parameter sets in $y = 0, 0.2$ and 0.5 , respectively.




Figure 2: The effective ω -meson mass m_ω^* is shown as a function of the baryon density ρ . The solid, dashed and dotted lines correspond to the results with the parameter sets for RHA, VS1 and VS2, respectively.